

Isolation and Identification of *Pseudomonas aruginosa* from Goldfish (*carassiusauratus*) and studing the antibiotic sensitivity in Al-Kufa city

Khalied Y.AL-Zamily

Ali Adnan AL- DarweshShoob N. Asal

Kut Tech. institute

Vet. College\ KufaVet.Hospital\Wassite

Abstract

This study was conducted to isolate and diagnose bacterial agents and studying the antibiotics sensitivity in gold fish (*carassiusauratus*). 43 goldfish were tested from the fish market in Al-Kufa city .The results of macroscopic examination showed hemorrhagic skin lesion , brown to red spotted skin of a varying degree were found as well as observed darkens color and the presence of black color on the fin. Also, noted the accumulation of fluid in the abdominal (ascitis) with no damage to internal organs observed. *Pseudomonasaruginosa* was isolated and diagnosed from the skin and the fins, Biochemical tests were conducted to confirm the diagnosis. The antibiotic sensitivity was tested by using seven antibiotics clindamycin (2 mg), ceftriaxone (30 mg), amikacin (30 mg), Sevbaudoxim (10 mg), ampicillin (10 mg), cefotaxime (30 mg), and gentamicin (10 mg)). The results of the sensitivity test were showed that amikacin, clindamycin and ceftriaxone was influential on the growth of bacteria with diameters Sense (30 ,25 and 20 mm) respectively, while remained antibiotics, the bacteria were resistant to them.

عزل وتشخيص جراثيم الزوائف النجارية من الاسماك الذهبية ودراسة حساسيتها للمضادات الحياتية في مدينة الكوفة

شعوب ناصر عسل

علي عدنان الدرويش

خالد ياسين الزامل

المستشفى البيطري\واسط

كلية الطب البيطري\كوفة

المعهد التقني\كوت

الخلاصة:

أجريت هذه الدراسة لعزل وتشخيص المسببات البكتيرية و دراسة حساسيتها للمضادات الحياتية من الأسماك الذهبية (القشري). تم فحص 43 سمكة ذهبية من سوق السمك في مدينة الكوفة. أظهرت نتائج الفحص العياني وجود آفات جلدية نزفية ولوحظت بقع حمراء مائلة للبني على الجلد بدرجات متفاوتة وكذلك لوحظ تغمق اللون و وجود اللون الأسود على الزعنفة، كذلك لوحظ تجمع السوائل في منطقة البطن مع عدم وجود اضطراب في الاحشاء الداخلية. تم عزل وتشخيص جراثيم *Pseudomonas aruginosa* من الجلد والزعانف وتم إجراء الفحوصات الكيموحيوية لتأكيد التشخيص. كذلك تم فحص مدى حساسيتها للمضادات الحياتية حيث استخدم سبعة مضادات حياتية (الكلينداميسين (2ملغ)، سيفترياكسون (30 ملغ)، الأميكاسين (30 ملغ، سيفيدوكسيم (10ملغ)، الأمبيسلين (10ملغ)، سيفوناكسيم (30ملغ)، وجنتاميسين (10ملغ). أظهرت نتائج فحص الحساسية ان الاميكاسين، الكلينداميسين و سيفترياكسون كانت مؤثرة على نمو البكتريا وباقطار تحسس (25,30 و 20 mm) على التوالي بينما بقيت المضادات الاخرى فكانت البكتريا مقاومة لها.

INTRODUCTION

Pseudomonas aruginosa are an ubiquitous group of environmental Gram-negative motile (have one flagella) (Stead., 1992). Bacterial pathogens in aquatic environments that can cause infections in fish, where a growing number of pathogenic bacteria causing the infection and death of fish (especially Koi fish freshwater aquaculture (Tor Monsenet *al.*, 2008). Several species of *Pseudomonas* have been reported cause disease in a number of fish species, include goldfish like *Carassiusauratus* (Bullock *al.*, 1965), and are

associated with septicemia in aquatic animals (Roberts *al.*, 1978). The lesions and hemorrhages were noted at the base of the fins or on the skin due to bacterial infection isolated bacteria were Gram negative, rod shaped and motile positive for oxidase, catalase, pyocyanin production, citrate utilization, gelatin liquefaction test. The bacteria were unable to ferment glucose. The biochemical properties and the characteristic blue-green appearance of culture due to the mixture of pyocyanin (blue) were indicative of *Pseudomonas aeruginosa*. (Somerita *et al.*, 2012). These bacteria have been considered as

opportunistic pathogens, causing diseases when the host is subjected to stress (Somsiri and Soontornvit ., 2002). The lesions, hemorrhages and ulceration were recorded at the base of the fins or on the skin due to bacterial infection (*P. aeruginosa*) (Hossain., 2008). This study aimed to determine the phenotypic and biochemical of the disease causing bacterial which isolated and determine effective antibiotic treatments.

MATERIAL AND METHOD:

1-Sampling of fish:

A total of 43 live fish were collected from fish farms located at NAJAF market. Afterward, the infected fishes were acclimatized in water of glass aquarium in the laboratory (University of Kufa, college of Veterinary medicine) conditions approximately at 25 °C.

2- isolation of Bacteria and culture:

Fish mucus, caudal fin and skin region were carefully scraped from the dorsal body using a sterilized cotton swab, ventral skin mucus was not collected to avoid intestinal and sperm contamination the sample were cultured in broth and agar incubated at 37 °C for 24 hour for growth and multiplication. Then the bacteria spread over the surface of MacConkey agar plate, Nutrient agar and pseudomonas agar. The pseudomonas agar was used for isolation of pure cultures of *Pseudomonas* species, the isolated bacteria were selected and characterized based on their biochemical properties (Holt and Krieg ., 1984). Dominant isolates were purified and identified using conventional biochemical tests described by (West and Colwell., 1984). API 20E test was done to sure the diagnosis.

Antibiotic sensitivity test:

Kirby-Bauer disk diffusion method was used to determine the antibiotic-resistant characteristics of the isolated organisms (Bauer et al., 1996). Study of resistance pattern of *Pseudomonas aeruginosa* isolates were cultured on the Muller-Hinton agar plate by using cotton swabs, and then 7 types of antibiotic were placed on the surface of agar by using sterile forceps

and incubated at 37 °C for 18 h. These antibiotics were Cefpodoxime (10 mg), Clindamycin (2 mg), Ampicillin (10 mg), Cefotaxime (30 mg), Gentamycin (10 mg), Ceftriaxone (30 mg), Amikacin (30 mg). The diameter of inhibition zones (mm) were recorded for all of the plates and then compared with the standard (National Committee., 1997).

Result and discussion

1-Clinical signs:

The clinical signs were shown. Lethargy and loss of appetite, gross lesions include black color on skin, fin is dark in color, the end of fin is damaged and necrotic (fig 1), ascites, internally the intestine was distended with clear, viscous fluid, hemorrhaging is common in the viscera and around the intestines, with enlargement of kidney, liver, and spleen and this agrees with (Somerita et al., 2012). The results also agreed with (Eissa et al., 2010) who stated ulceration and necrotic lesions of skin, fin and gill rot, accumulation of abdominal fluid, haemorrhagic septicemia and fish mortality. The lesions and hemorrhages were noted at the base of the fins or on the skin due to bacterial infection. Fluid accumulation in the abdomen is a common feature; the liver and spleen were distended and fluid filled.

2- Organismic Identity:

P. aeruginosa Colony on the nutrient agar was blue green fluorescent (fig 2), on MacConkey agar the colonies were blue green pigment because the *Pseudomonas aeruginosa* produces lactose non-fermentation and this agrees with (Tor Monsen et al., 2008), (fig 3). Also these results agreed with (Akinyemi., 2012) who isolated *P. aeruginosa* from the gill, skin. Also our results agreed with those of (Hossain et al., 2006; Musa et al., 2009). Biochemical test delivered a preliminary identification of the *Pseudomonas aeruginosa* strain. *Pseudomonas aeruginosa* isolates were identified as Gram negative, motile, oxidase-positive will show a blue color due to produce the enzyme cytochrome-c oxidase, and ability to reduce oxygen and this consent with (West et al., 1985). Simmons

citrate-negative will show green mean no. growth because unable to utilize citrate as the sole carbon and energy source and this assent with (Collins and Lawson2000) and urease negative because there is no rise in pH due to accumulation of ammonia and this approved with (Christensen.,1946). *P.aureginosa* were parallel to previous studies which identified the same organisms from other fish species (Austin and Austin 2007). *P. aeruginosa* are opportunistic Gram negative pathogens, naturally occurring in aquatic environment and as a part of normal gut flora of healthy fish, it cause outbreak when the normal environmental conditions changed (Angelini and Seigneur.,1988).

3- Antibiotic susceptibility

The antibacterial activity of seven drugs type were evaluated bacterial isolates *P.aeruginosa*(fig 3) Which exhibited more sensitive with tested antibacterial drugs in Clindamycin(2mg), Ceftriaxone(30mg) and Amikacin(30mg) the inhibition zones of other antibacterial drugs were (30,25,20 mm) respectively and the *Pseudomonas aeruginosa* was resistant with Cefpodoxime(10 mg), Ampicillin(10mg), Cefotaxime(30mg) and Gentamycin(10mg),(Table 1)The variation in

antibiotic sensitivity test of isolated pathogen was due to *P. aeruginosa* is intrinsically resistant to several antibiotics because of the low permeability of its outer-membrane, the constitutive expression of various efflux pumps, and the production of antibiotic-inactivating enzymes

(e.g.,cephalosporinases)(Hancock.,1998).Also these variation may be related to the large size and the versatility of its genome, and to its distribution in aquatic habitats, which could constitute a reservoir for bacteria carrying other resistance genes (Vaisvila*et al.*,2001).Also variation may be depend on culture circumstances and highlight(Dominic *et al.*,2005).This result agreed with (Saraswathi *et al.*,2013).(Algama*et al.*,2012)who showed the *P. aeruginosa* sensitive to Gentamycin these results disagreement with our results, These isolates were high resistance against Kanamycin, Cefpodoxime, Cephodoxime. Also, (Eissa*et al.*,2010)who experimented the antibiogram sensitivity ,he observed *P.* highly sensitive to Amikacin and sensitive to Gentamicin. We concluded that *P. aeruginosa* isolates were identified the most common bacteria isolated from gold fish, *P. aeruginosa* isolates were sensitive to Clindamycin, Ceftriaxone and Amikacin.

Table (1) Inhibition zones of seven antibiotic drugs against *Pseudomonas aeruginosa* in vitro.

Types and concentration of antibiotic	Inhibition zone (mm)
Clindamycin(2mg)	25±1.9
Ceftriaxone(30mg)	20±2.1
Amikacin(30mg)	30±2.30
Cefpodoxime(10 mg)	-
Ampicillin(10mg)	-
Cefotaxime(30mg)	-
Gentamycin(10mg)	-



Fig 1 . Gross lesion include black color on skin and fin




Fig 2: *P.aeruginosa* Colony on MacConkey agar




Fig 3: Inhibition zones of seven antibiotic drugs against *Pseudomonas aeruginosa*

REFERENCE:

- Akinyemi, A. A (2012) : Plant extracts as alternative treatment for *Pseudomonas aeruginosa* occurrence in *Clarias gariepinus* (Burchell, 1822) juveniles. Journal of Science and Multidisciplinary Research : 39-50.
- Algama D.W.R. R ; Kurcheti P.P.; Subhas C.M. ; Kumar K., Rajeev K. B., Charandas T. M. and Neeraj K.(2012) : *In vitro* Sensitivity of Three Bacterial Pathogens of Koi Carp (*Cyprinus carpio* L.) to Certain Antibiotics. Journal of Agricultural Science and Technology B 2 :93-98.
- Angelini, N.M., and Seigneur, G.N., 1988. Disease of the fins of *Rhamdia sapo*. Isolation of the etiological agents and experimental infection. Rev. Argent. Microbiol. 20, 37– 48.
- Austin, B. and Austin, D. A. 2007, Bacterial Fish Pathogens, Diseases of Farmed and Wild Fish. Fourth Edition, Praxis Publishing Ltd, Chichester, UK.
- Bauer, A.W.; Kirby, M.M.; Sherris, J.S. and Tenckhoff, M. (1996). Antibiotic sensitivity testing by single disc method. Am. J. Clin. Pathol. (45): 939-396.
- Bullock, G.L. (1965). Characteristics and pathogenicity of a capsulated *Pseudomonas* isolated from goldfish. Appl. Microbiol. 13:89-92.
- Christensen, W.B. (1946) Urea decomposition as a means of differentiating *Proteus* and paracolon cultures from each other and from *Salmonella* and *Shigella* types. Journal of Bacteriology 52, 461–466.
- Collins, M.D. and Lawson, P.A. (2000) .The genus *Abiotrophia* (Kawamura *et al.*) is not monophyletic; proposal of *Granulicatella* nov., *Granulicatella adiacens* comb. nov., *Granulicatella elegans* comb. nov., and *Granulicatella balaenopterae*. International Journal of Systematic and Evolutionary Microbiology 50, 365–369.
- Dominic, H., Barbara, R., Aniko, P., Michael, R., Peter, B., Scott, B., Mark, E., Barbara, T., Colin, M., Shawn, D.A. and Colin, H. 2005. Antibiotic Susceptibilities of *Pseudomonas aeruginosa* Isolates Derived from Patients with Cystic Fibrosis under Aerobic, Anaerobic, and Biofilm Conditions. J. Clin. Microbiol. 43 (10): 5085-5090.
- Eissa. N.M.E. , Abou El-Ghiet. E.N. , A.A. Shaheen. A.A. and Abbass. A. 2010:

- Characterization of *Pseudomonas* Species Isolated from Tilapia "*Oreochromis niloticus*" in Qaroun and Wadi-El-Rayan Lakes, Egypt . Global Veterinaria 5 (2): 116-121.
- Hancock ,R.E.(1998). Resistance mechanisms in *Pseudomonas aeruginosa* and other non fermentative gram-negative bacteria. Clin Infect Dis ; 27 (suppl 1): S93–S99.
- Holt, J.G. and Krieg, N.R., 1984. Bergey's Manual of Systematic Bacteriology vol. 1. Williams and Wilkins, Baltimore, USA.
- Hossain M.A. (2008). Isolation of pathogenic bacteria from the skin ulcerous symptomatic gourami (*Colisa lalia*) through 16S rDNA analysis. Univ. J. Zool. Rajshahi Univ. pp. 21-24.
- Hossain,M.I., Farzana A.N., Hussain,M.A., Rahman,M.H and Satoru, S. (2006) : Disribution of *Pseudomonas aeruginosa* in swamps and its infection to *Oreochromis niloticus*. J. bio-sci. 14: 77-81.
- Musa,N., Wei, L.S. and Wendy, W. (2009): Bacterial diseases outbreak of African Catfish (*Clarias gariepinus*) from Manir River, Terengganu, Malaysia. J lif. Sci.,volume 3 no (5) :10-20.
- National Committee for Clinical Laboratory Standards.(1997). Performance standards for antimicrobial disk susceptibility tests. Approved standard M2-A6. Wayne, Pa: National Committee for Clinical Laboratory Standards.
- Roberts ,R.J. (1978). Fish Pathology 2nd ed. Bailliere Tindall, London. Saitou N, Nei M (1987). The neighbor-joining method: a new method for reconstructing phylogenetic trees. Mol. Biol. Evol. 4:406-425.
- Saraswathi N.K. Ashadevi, N. Kanipandian, 1 1 2V. Vignesh and 2R. Thirumurugan.(2013).Studies on Antibiotic Resistance Pathogens Isolated from Infected Ornamental Fish World Journal of Fish and Marine Sciences 5 (5): 540-544.
- Somerita P., P. K. Bandyopadhyay¹ and S. N. Chatterjee (2012). Characterization of *Pseudomonas aeruginosa* PB112 (JN996498) isolated from infected *Labeo bata* (Hamilton) by 16S rRNA gene sequence analysis and fatty acid methyl ester (FAME) analysis. pp. 400-405, 23.
- Somsiri T. and Soontornvit S. (2002). Bacterial diseases of cultured tiger frog (*Rana tigrina*). Diseases in Asian Aquaculture IV, Fish Health Section, Asian Fisheries Society, Manila. pp. 15-17.
- Stead, D., 1992. Grouping of plant pathogens and some other *Pseudomonas spp.* By using cellular fatty acid profiles. Int. J. Syst. Bacteriol. 42, 281–295.
- Tor Monsen, Elisabert Lovgren Micael Widerstrom and Lars Wallinder, 2008. In Vitro Effect of Ultrasonic on Bacteria and Suggested Protocol for Sonication and Diagnosis of Prosthetic Infection, University Hospital of Umeå Sweden, Journal of Clinical Microbiology.
- Vaisvila R, Morgan RD, Posfai J, Raleigh EA.(2001). Discovery and distribution of super-integrins among pseudomonads. Mol Microbiol; 42: 587–601.
- West, P.A. and Colwell, R.R., 1984. Identification of Pseudomonadaceae: an overview. In: Colwell, R.R. (Ed.), Pseudomoniasis in the Environment. John Wiley, New York, USA, pp. 141– 199.
- West, P.A., Brayton, P.R., Twilley, R.R., Bryant T.N. and Colwell, R.R. (1985) Numerical taxonomy of nitrogen fixing 'decarboxylase-negative' *Vibrio* species isolated from aquatic environments. International Journal of Systematic Bacteriology 35, 198–205.

