

## Detection of intracellular adhesion (*ica*) B gene in *Staphylococcus aureus* isolated from wound infections

Abbas D. Mater<sup>2</sup>, Mohammed Jasim Mohammed Shallal<sup>1</sup>([mohammed-j@utq.edu.iq](mailto:mohammed-j@utq.edu.iq)),  
Zainab Dakhil Degaim<sup>1</sup>([zainab-d@utq.edu.iq](mailto:zainab-d@utq.edu.iq))

<sup>2</sup>Basic Sciences department-College of Dentistry / University of Thi-Qar, Iraq.

<sup>1</sup>Microbiology Department-College of Medicine-University of Thi-Qar, Iraq

### Abstract

**Background:** *Staphylococcus aureus* is an imperative cause of community acquired infections and nosocomial infections, The formation of biofilm on host surfaces and adherence is painstaking to be significant virulence factor in *S. aureus*.

**Objective:** The main objective of this study is to detect and describe the distribution of *S. aureus* and *icaB* gene in wound infection samples taken from patients at Al-Hussein Teaching Hospital in Thi-Qar province, Iraq, during the period extended from February to September 2007.

**Materials and Methods:** A total number of bacterial isolates included 130 sample recovered from different wounds of patients in burn unit. All isolates of the targeted bacteria were subjected for screening the biofilm allied gene (*icaB*) by Polymerase Chain Reaction (PCR) technique.

**Results:** The recent data indicated that 72 (55.3%) identified as *S. aureus*, and the females were likely more affected with *S. aureus* infection than males (47 > 25). The results of PCR recorded that *icaB* gene was expressed only in 61% of all isolates.

**Conclusion:** The recent results recorded a high percentage of *icaB* gene which related with biofilm production may be increased the pathogenicity of this pathogen which caused different human diseases.

**Keywords:** *S. aureus*, PCR, *icaB*.

## Introduction:

*Staphylococcus aureus* is an imperative cause of community acquired infections and nosocomial infections, and it caused a surfeit of diseases like pneumonia, skin, soft tissue infections, and blood stream infections<sup>1</sup>. One of the important virulence factors of *S. aureus* is a strain of staphylococci characterized by different levels of virulence. The virulence of this type of bacteria is recognized by its ability to form biofilms. Biofilm is a functional factor necessary for the attachment of these bacteria to the surfaces and their embedding in various extracellular polymeric substances. Intracellular adhesion locus (*ica*) is considered as a group of final biofilm product of that produced by *S. aureus* which involves complex mechanism<sup>2</sup>.

The formation of biofilm on host surfaces and adherence is painstaking to be significant virulence factor in *S. aureus*<sup>3</sup> that controlled by various virulence factors expressed during *Staph. aureus* nosocomial infections and also by genes encoding antibiotic resistance that frequently presented. Cramton *et al.*<sup>3</sup> documented that the intracellular adhesion locus (*ica*) is required for biofilm production, also whom

suggest that the *ica* locus could potentially be an important target in the therapy of implant infections<sup>4</sup>, and approximately 60% of *S. aureus* strains were produced the biofilm<sup>5</sup>. While, contradictory the study performed by Rohde *et al.*<sup>6</sup> suggested that all *S. aureus* strains possess *icaADBC* genes. The first identification of *icaADBC* operon in *Staph. epidermidis* followed by showing to be current in *S. aureus*<sup>7</sup>. Although Arciola *et al.*<sup>8</sup> reported the greatest of *S. aureus* isolates appear to contain this operon, and *icaADBC* operon is consist of the four genes including: *icaA*, *icaB*, *icaC* and *icaD*<sup>9</sup>. The formation of biofilm is influenced by amount of factors, and the synthesis of the polysaccharide intercellular adhesion (PIA) by the bacteria is the most important factors that increase the bacterial virulence in addition to their support to the survival of *Staph. aureus* in a variety of environments<sup>7</sup>. Also *icaADBC* genes encodes for the synthesis of PIA<sup>10</sup>. A *icaB* is the deacetylase responsible for the de-acetylation of mature PIA and the trans-membrane protein<sup>11</sup>. The operon expression augmented under anaerobic internal environment of biofilm<sup>12</sup>. The aim of this study was to investigate the frequency of the biofilm producing related gene (*icaB*) in *S.*

*aureus* isolates originated from wound infections.

### Material and Methods:

**Ethical approval** This research was approved by the Medical College Ethics Committee, Thi-Qar University, Thi-Qar Province, Iraq.

**Type of study:** Cross sectional descriptive study extend all over February to September 2007. In total, 130 bacterial isolates were taken from patients hospitalized in the burn unit at Al-Hussein Teaching Hospital in Thi-Qar from February to September 2007. All specimens were taken from wounds and the bacteria were maintained in a sterile transport medium (Amies transport medium) used for bacterial isolation.

### Laboratory methods

All strains of *S. aureus* were isolated from 130 swabs which collected from burn wound infections identified depending on cultural properties on different culturing media, followed by biochemical tests<sup>13</sup>. The confirmed diagnosis was performed by using Analytical Profile Index system Staph (API) (BioMerieux/France).

### Preparation of bacterial DNA

Genomic DNA was extracted from all isolates by using Genomic DNA Extraction kit (Geneaid/Korea).

### PCR Protocol

All isolates of *S. aureus* (72/130) were subjected to amplification of *icaB* gene. The specific primer pairs of *icaB* are following: forward: 5'-CCC AAC GCT AAA ATC ATC GC-3' and reverse: 5'-ATT GGA GTT CGG AGT GAC TGC-3'. The PCR cycling conditions of current gene: initial denaturation at 94°C for 5 min, followed by 30 cycles of denaturation at 94°C for 1 min, annealing at 53°C for 1 min, extension at 72°C for 1 min and final extension for 9 min<sup>14</sup>.


### Results

Among 130 of bacterial isolates taken from wound infections of burn patients, only 72 (55.38%) were positive for *S. aureus* recovered from all bacterial isolates. Also, it was detected that only 47 out of 71 bacterial isolates was positive for *S. aureus* in female with a percentage of was 65.3%. However, this was less in males, 25 out of 59 with a percentage of 34.7% as shown in table 1.

The PCR results of *ica B* gene were detected and recognized by agarose gel electrophoresis that measured by bands of size 1080 bp as presented in figure 1.

**Table (1):** Percentage of *S. aureus* isolated recovered from bacterial isolates taken from wounds of patients of burn infection among males and females.

Gender	No	No. of <i>S. aureus</i> (+ve)
Male	59	25 (34.7%)
Female	71	47 (65.3%)
Total	130	72



**Fig. (1):** Agarose gel electrophoresis of *ica B* gene amplification ligated PCR used a 1.4% agarose gel, Lane M: DNA ladder Lanes, 1-8: These bands of size 1080 bp represent the positive results of amplified *ica B* gene

**Discussion:**

The results of the recent study is showed that 72 of bacterial isolates (55.38 %) were characterized as *S. aureus* from all collected samples. The *S. aureus* is one of the most significant causes of wound infections. Fitzpatrick *et al.*<sup>15</sup> recorded that *staphylococcus* spp. which produced the biofilm had become more predominant as cause of the hospital-acquired infection. The results of current study are analogous with results of Alwashand Saleh<sup>23</sup>, described the occurrence of *S. aureus* in burns was 33.3%.

The results of present study documented that the females were likely more affected with *S. aureus* infection than males (47 >

25). The current results were in agreement with the results of Rajput *et al.*,<sup>25</sup> indicated that burn infection in females was (60%). While the same results of the present work were dissimilar to study that conducted in Iraq by Abid,<sup>24</sup> showed that males tend to be more infected with *S. aureus* than women with 52.2%.

The occurrence of the biofilm allied gene (*icaB*) amongst entirely a *S. aureus* isolates showed that *icaB* gene was expressed in 61% of the bacterial isolates. The adherence on surfaces and biofilm formation on wounds is associated with increased the pathogenesis of *S. aureus* according to previous studies which proved this truth<sup>16</sup>, and the biofilms help microbes to survive

hostile environments such as antibiotics and the host immune response<sup>16,17</sup>. Also the initial detection of biofilm formation by *Staphylococcus* must be the crucial steps in the direction of the prevention of nosocomial infections<sup>18</sup>.

*S. aureus* had the ability to produce the biofilm on an inert or living surface<sup>19,20</sup>; its capacity which supports this pathogen for causing the infection and severe morbidity<sup>21</sup>.

The current results incorporated with results of the study performed by Mirzaee et al.<sup>22</sup> recorded that the prevalence of *icaB* gene in Methicillin resistant *S. aureus* (MRSA) isolates was 51%.

### Conclusion:

The recent data recorded a high percentage of *icaB* gene which associated with biofilm production may be increased the pathogenicity of this pathogen which caused different human diseases.

### Reference

[1] Havaei, S.; Azimian, A.; Fazeli, H.; Naderi, M.; Ghazvini, K.; Samiee, S M.; Masoumi, Z. and Akbari, M. (2012). Genetic characterization of Methicillin Resistant and Sensitive, Vancomycin Intermediate *Staphylococcus aureus* strains isolated from different Iranian hospitals. J. ISRN. Microbiol., 20:215275.  
[2] Shojaei, M.; KarimiDarehabi, H. and Javadi, A. (2014). Evaluation of the

prevalence of genes producing biofilm (*fnbB*, *clfaA*, *icaC*, *icaB*) in *S.aureus* strain isolated from raw milk in sanandaj province. J. Food Process Technol. 5:4.

[3] Cramton, S E.; Gerke, Ch.; Schnell, N F.; Nichols, W W. and Gotz, F. (1999). The intercellular adhesion (*ica*) locus is present in *Staphylococcus aureus* and is required for biofilm formation. J. Infect. Immun., 67: 5427–5433.

[4] Arciola, C R.; Baldassarri, L. and Montanaro, L. (2001a). Presence of *icaA* and *icaD* genes and slime production in a collection of staphylococcal strains from catheter-associated infections. J. Clin. Microbiol., 39:2151-2156.

[5] Arciola, C R.; Campoccia, D.; Borelli, AM. and Montanaro, ME. (2001b). Congo red agar plate method: improved accuracy and new extended application to *Staphylococcus aureus*. J. Microbiol., 24: 355–363.

[6] Rohde, H.; Knobloch, JK.; Horstkotte, M A. and Mack, D. (2001) Correlation of *Staphylococcus aureus* *icaADBC* genotype and biofilm expression phenotype. J. Clin. Microbiol., 39: 4595–4596.

[7] Gotz, F. (2002). *Staphylococcus* and biofilms. J. Mol. Microbiol., 43:1367-1378.

[8] Arciola, C R.; Campoccia, D.; Ravaioli, S. and Montanaro, L. (2015). Polysaccharide intercellular adhesin in

biofilm: structural and regulatory aspect. J. Front. Cell. Infect. Microbiol., 5:1-10.

[9] O’Gara, J P. (2007). *ica* and beyond: biofilm mechanisms and regulations in *Staphylococcusepidermidis* and *Staphylococcus aureus*. J. FEMS. Microbiol., 270: 179-188.

[10] Tiemersma, E W.; Bronzwear, L A.; Lyytikäinen, O.; Degener, J E.; Schrijnemakers, P.; Bruinsma, N.; Monen, J.; Witte, W. and Grundman, H. (2004). Methicillin resistant *Staphylococcus aureus* in Europe, 1999-2002. J. Emerg. Infect. Dis., 10:1627-16134.

[11] Gamal, F.; Mohamed, A.; Mostafa, S.; Mona, A.; Hassan, A. and Rehab, M. (2009). Detection of *icaA*, *icaD* genes and biofilm production by *Staphylococcus aureus* and *Staphylococcus epidermidis* isolated from urinary tract catheterized patients. J. Infect. Dev. Ctries., 3 (5):342-351.

[12] Glowalla, E.; Tosetti, B.; Kronke, M. and Krut, O. (2009). Proteomics based identification of anchorless cell wall proteins as vaccine candidates against *Staphylococcus aureus*. J. Infect. Immun., 77 (7):2719-2729.

[13] Harley, J P. and Prescott, L M. (2002). Laboratory Exercises in Microbiology. (5<sup>th</sup>ed). The McGraw-Hill Companies, Inc., New York.

[14] Kim, J H. ; Kim, C H.; Hacker, J.; Ziebuhr, W.; Lee, B K. and Cho, S H.(2008). Molecular characterization of regulatory genes associated with biofilm variation in a *Staphylococcus aureus* strain. J. Micro. Biotech. 18(1): 28-34.

[15] Fitzpatrick, F.; Humphreys, H.; Smyth, E.; Kennedy, C A. and O’Gara, GP. (2002). Environmental regulation of biofilm formation in intensive care unit isolates of *Staphylococcus epidermidis*. J. Hosp. Infec., 42:212-218.

[16] Grinholc, M.; Wegrzyn, G. and Kurlenda, J. (2007). Evaluation of biofilm production and prevalence of the *icaD* gene in methicillin-resistant and methicillin-susceptible *Staphylococcus aureus* strains isolated from patients with nosocomial infections and carriers. J. FEMS Immune. Med. Microbiol., 50:375-379.

[17] Eftekhari, F. and Dadaei, T. (2011). Biofilm formation and detection of *icaAB* genes in clinical isolates of Methicillin Resistant *Staphylococcus aureus*. Irani. J. Basi. Med. Scien., 14(2):132-136.

[18] Nasr, R A.; Abu-Shady, H M. and Hussein, H S. (2012). Biofilm formation and presence of *icaAD* gene in clinical isolates of *staphylococci*. Egyp. J. Med. Hum. Genet., 13:269-274.

- [19] Costerton, J W.; Stewart, P S. and Greenberg, E P. (1999). Bacterial biofilms: a common cause of persistent infections. *J. Science*. 284 (5418): 1318 – 1322.
- [20] Dhanawade, N B.; Kalorey, D R.; Srinivasan, R.; Barbuddhe, S B. and Kurkure, N V. (2010). Detection of intercellular adhesion genes and biofilm production in *Staphylococcus aureus* isolated from bovine subclinical mastitis. *J. Vet. Res. Commun.*, 34: 81-89.
- [21]Waldvogel, FA. (1995). In: principles and practices of infectious diseases. Mandell, Douglas, and Bennett ed. Wiley. New York. pp 1754-1777.
- [22] Mirzaee, M.; Peerayeh, S N. and Ghasemian, A. (2014). Detection of *ica*ABCDgenes and biofilm formation in clinical isolates of Methicillin Resistant *Staphylococcus aureus*. *Iranin. J. Patho.*, 9 (4):257 – 262.
- [23] Alwash , S A. and Saleh, D S. (2013). Comparison between Cefoxitin disk diffusion, Crome agar and EPI-M screening kit for detection of Methicillin Resistant *Staphylococcus aureus*. *Iraqi. J. Sci.*, 54(4):847-850.
- [24] Abid, A N. (2015). Serological diagnosis and Molecular detection of Methicillin resistant *Staphylococcus aureus* isolated from burn patients in Al-Hussein Teching Hospital at Thi-Qar Province. M.Sc. Thesis. College of Science/ Thi-Qar University. Iraq.
- [25] Rajput, A.; Singh, K.P.; Kumar, V.; Sexena, R. and Singh, R K. (2008). Antibacterial resistance pattern of aerobic bacteria isolates from burn patients in tertiary care hospital. *J. Biomed. Res.*, 19(1): 1-4.

## الكشف عن وجود جين الالتصاق الخلوي في المكورات العنقودية الذهبية المعزولة من أصابات الجروح

د. عباس دخيل مطر<sup>٢</sup>

د. محمد جاسم محمد شلال<sup>١</sup>

د. زينب داخل دغيم<sup>١</sup>

<sup>١</sup> فرع الأحياء المجهرية-كلية الطب- جامعة ذي قار

<sup>٢</sup> قسم العلوم الأساسية- كلية طب الأسنان- جامعة ذي قار

### الملخص:

**الهدف:** تعتبر بكتريا المكورات العنقودية الذهبية احد المسببات الاساسية لحالات الاصابة في المجتمع بالإضافة الى العدوى الناتجة في المستشفيات. ان تكوين biofilm على سطوح خلايا المضيف والتي تساعد على التصاق هذه الاحياء المجهرية على سطوح تلك الخلايا هي احد عوامل الضراوة المهمة للمكورات العنقودية الذهبية.

**المواد وطرق العمل:** من مجموع ١٣٠ عينة من المكورات العنقودية الذهبية تم جمعها الجروح في مستشفى الحسين التعليمي في محافظة ذي قار، العراق. حيث تم استخدام كل عزلات البكتريا الهدف للكشف عن الجين المسؤول لتكوين biofilm (*icaB*) بواسطة تقنية تفاعل سلسلة البلمرة الجزيئية (PCR).



**النتائج:** بينت النتائج الحالية ان ٧٢ (٥٥.٣%) تم تشخيصها على انها بكتريا المكورات العنقودية الذهبية. كما اوضحت النتائج الجزيئية لاختبار تفاعل سلسلة البلمرة الجزيئية للجين الهدف ان ٥١.١٥% من العزلات تحتوي على هذا الجين.

الاستنتاج: اوضحت النتائج الحالية نسبة عالية من وجود جين *icaB* ذات العلاقة بتكوين biofilm والذي قد يكون لع علاقة بزيادة الامراضية لهذه البكتريا المسببة لانواع متعددة من الامراض للانسان.