

THE INFLUENCE OF HEAVY METALS AND ANTIMICROBIAL ON STAPHYLOCOCCUS AUREUS AND PSEUDOMONAS AEROGINOSA ISOLATES

Mohammed H. Khudor

Department of Microbiology, College of Veterinary Medicine, University of Basrah-
Basrah-Iraq.

(Received 7 September 2010 , Accepted 20 October 2010)

Keywords; *Staphylococcus aureus* , gentamycin, chloramphenicol

ABSTRACT

Two isolates of *Staphylococcus aureus* and *Pseudomonas aeruginosa* of bacteria were exposed to different concentrations (5 , 10 or 25 mg /l)of heavy metals ions (Silver Ag^{2+} , Cobalt Co^{2+} or Lead Pb^{2+}) for different exposure periods (0 , 1, 2, 24 and 48 hrs.) .The results showed that the inhibitory effect of these metals on the growth of the two isolates were in the following sequences :- $Pb < Co < Ag$

Some of these metals caused an elongation of lag phase in liquid medium for these isolates . When antibiotic discs (chloramphenicol 30 mcg , clindamycin 2 mcg , erythromycin 15 mcg , gentamycin 10 mcg , tetracyclin 30 mcg and vancomycin 30 mcg) were added to solid medium which was previously supplemented with the above heavy metals ions , a synergistic effect was observed between antibiotics and heavy metals ions (specially silver ions at 25 mg/ l)to increase the inhibition zone significantly ($P < 0.01$) for both isolates .

INTRODUCTION

The high frequency of antibiotic resistant staphylococci , even among those not known to have been treated with antibiotics would obscure subtle changes in the microbial flora (1). Multidrug resistance strains of *Pseudomonas aeruginosa* are the major cause of mortality in various clinical conditions , such as cystic fibrosis (2 , 3 , 4 , and 5) and burn-wound infections (6 , 7 and 8) especially in nosocomial situations . This condition has led to the development of new or modified antimicrobial agents

for the therapy of such infections . Heavy metals , particularly silver and mercury , have a variety of applications in controlling microbial populations (9) .

Microorganisms require some metals like Cu^{2+} , Zn^{2+} , Co^{2+} and Ni^{2+} at low concentrations as essential micronutrients for vital cofactors for metalloproteins and certain enzymes , however , at high concentration it has been reported that these metals interact with nucleic acids and enzyme active sites. Toxic effects include ion displacement and / or substitution of essential ions from cellular sites and blocking of functional groups of important molecules , e.g. , enzymes , polypeptides , and essential nutrients transport system (10) .

Silver salts alone or in combination with other drugs appear to have a significant potential as antimicrobial agents (11 , 12 , and 13) . Similarly , mercury in the form of less toxic organic compounds is being used as skin disinfection (14) .

Some novel transition metal (Cu^{2+} , Zn^{2+} , Co^{2+} and Ni^{2+}) complexes of substituted pyridine Schiff- bases have been prepared in order to evaluate the effect of metal ions upon chelation , the Schiff- bases and their complexes as antibacterial against *Escherichia coli* , *Staphylococcus aureus* and *Pseudomonas aeruginosa* , the complexed Schiff- bases have shown to be more antibacterial against one more bacterial species as compared to uncomplexed Schiff- bases (15) .

Ahmed et al (16) synthesized complexes of Co^{2+} , Ni^{2+} , Cu^{2+} , Mn^{2+} , Zn^{2+} , Fe^{2+} , Fe^{3+} and La^{3+} with methylenedisalicylic acid (MDSA) and revealed a high antimicrobial activity for *S. aureus* and *P. aeruginosa* . Very little information is available in the literature regarding the action of heavy metals against clinical isolates (17) .

This study was carried out to assess the effects of heavy metal ions on *S. aureus* and *P. aeruginosa* , and to assess the effect of compensation between heavy metals and antibiotics on the above microorganisms.

MATERIALS AND METHODS

Two isolates *Staphylococcus aureus* and *Pseudomonas aeruginosa* were obtained from a bacteriological lab. at College of Science, University of Basrah . Nutrient Agar (NA) was used for culturing and activation of these isolates for 24 hrs. , then bacterial cell suspension was prepared in the concentration 1×10^6 cell / ml by using Petroff-Hausser counting chamber(18) . One ml of this cell suspension was added to each

tube which containing 20 ml of nutrient broth (NB) medium supplemented with one concentration (5 , 10 or 25 mg/ l)of heavy metals ions : Ag as AgNO₃ ,pb as PbNO₃ or Co as Co(NO₃)₂ .6H₂O) , triplicates were done for each treatment in addition to control . These cultures were incubated at 37 °C . The growth rate (OD at 600 nm) was measured after different exposure periods (0 , 1 , 2 , 24 or 48 hrs.) . From each treatment 0.2 ml was streaked (after 24 hrs. exposed period) on Muller – Hinton Agar (MHA) medium free of metals ions , left for 15 min. to dry . Commercial antibiotics discs(4 discs / plate) were used (Table 1) .

The diameter of inhibition zones were recorded after overnight incubation at 37° C.

Another experiment for antibiotic sensitivity test was done on a defined medium (1.5 % w/ v MHA) using plates containing solidified medium (15 ml / plate) supplemented with heavy metals ions . A sample for a late - exponential phase culture(0.3 ml) containing about 1×10⁶ cell / ml was streaking onto the plates and allowed to dry . Commercial antibiotics discs were used at the concentrations that mentioned in table 1. The diameter of the inhibition zones were recorded after overnight incubation at 37 °C .

Data were analyzed statistically by using the analysis of variance (ANOVA test) and revised least significant differences test (RLSD) to compare the means .

Table (1) : Interpretation of inhibition zone diameter (Bioanalyse sensitivity discs Ankara / Turkey) .

Antibiotic	Symbol	Concentration	Diameter of inhibition zone	
			Resistant	Sensitive
Chloramphenicol	C	30 mcg	≤ 12	≥ 18
Clindamycin	DA	2 mcg	≤ 14	≥ 17
Erythromycin	E	15 mcg	≤ 13	≥ 18
Gentamycin	CN	10 mcg	≤ 12	≥ 15
Tetracyclin	TE	30 mcg	≤ 14	≥ 19
Vancomycin	VA	30 mcg	≤ 9	≥ 12

RESULTS AND DISCUSSION

The effect of heavy metals ions on the growth of *P. aeruginosa* in the liquid medium.

1- The effect of silver ions (Ag^{1+}):-

The exposure of bacteria to silver ions at 5 or 10 mg/l concentrations lead to increase the lag phase period comparable with the control treatment , while the concentration 25 mg / l lead to significant decreasing ($p < 0.01$) in the growth of bacteria without increasing in the lag phase period(Fig. 1)




Fig (1) : Effect of silver ions on the growth of *Pseudomonas aeruginosa* in the liquid medium at different exposure periods.

2- The effect of cobalt ions (Co^{2+}) :-

The effect of cobalt ions at 5 mg / l concentration was not differ significantly comparable with the control treatment . On the other hand ,10 mg / l concentration lead to relative decreases in the growth comparable with control treatment , while 25mg / l of cobalt ions lead to elongation the lag phase more than 24 hrs., with relative decreases in the growth rate (Fig. 2).

3- The effect of lead ions (Pb^{2+}) :-

The growth of bacteria was not differ significantly comparable with the control treatment at all concentrations of lead ions , and at all the period of the experiment (Fig. 3).




Fig (2) : Effect of cobalt ions on the growth of *Pseudomonas aeruginosa* in the liquid medium at different exposure periods.




Fig (3) : Effect of lead ions on the growth of *Pseudomonas aeruginosa* in the liquid medium at different exposure periods.

The effect of heavy metals ions on the growth of *S. aureus* in the liquid medium .

1- The effect of silver ions (Ag¹⁺):-

Silver ions at 5 mg /l was elongated the lag phase period more than 24 hrs. , and caused decreasing the growth after 48 hrs. While 10 mg /l revealed significant decreasing in the growth of bacteria without increasing in the lag phase periods , but 25 mg /l caused decreasing in the growth rate after 48 hrs. with increasing the lag phase for more than 24 hrs. (Fig. 4).




Fig (4) : Effect of silver ions on the growth of *Staphylococcus aureus* in the liquid medium at different exposure periods.

2- The effect of cobalt ions (Co ²⁺) :-

The growth rate of bacteria was decreased at 5mg /l concentration without an effect on the lag phase period . Ten mg /l concentration was elongated the lag phase period and decreased the growth rate , while 25 mg /l concentration was decreased the growth after 1 and 2 hrs. of exposure period ., and no growth was appeared after 24 or 48 hrs. of the exposure period (Fig. 5).

3- The effect of lead ions (Pb²⁺) :-

The growth rate was decreased when lead was used at 5 , 10 or 25 mg /l concentrations , without an effect on the lag phase period . The growth rate was significantly decreased ($p < 0.01$) in comparable with the control treatment (Fig. 6)




Fig (5) : Effect of cobalt ions on the growth of *Staphylococcus aureus* in the liquid medium at different exposure periods.




Fig (6) : Effect of lead ions on the growth of *Staphylococcus aureus* in the liquid medium at different exposure periods.

The different effect probably depended on the strain and on the experimental conditions such as: strain source , time of exposure (19), and type of element , the mechanism of efflux system that differ between *S. aureus* and *P. aeruginosa* (10).

Also the difference depended on the differences in the wall binding sites to interact with the metal ions , and differ in metal ions affinity to interact with the wall binding sites (20).

Pseudomonas aeruginosa exhibited increasing lag phase in growth after exposure to high concentrations of the heavy metals .This observation appears to be due to the selection for metal resistant phenotype (10).

The synergistic effect of heavy metals ions and antibiotics on the growth of *P. aeruginosa* in the solid medium

To investigate the co-effect of heavy metals with antibiotics on the tested bacteria . The results of experiment was revealed significant increasing ($P < 0.01$) in the inhibition zone when used chloramphenicol antibiotic with silver or cobalt ions in the solid medium plates (1,2) , and no growth appeared at chloramphenicol antibiotic with 25 mg/l silver ions . On the other hand , the inhibition zone do not differ significantly ($P > 0.05$) in treatments which containing chloramphenicol with lead ions, plate(3) comparable with the control treatment which was free from any addition of metals ions, table (2) .

Table (2) : Inhibition zone diameter (mm) of antibiotic alon and with heavy metals ions on *P. aeruginosa* at the solid medium

Antibiotic	Alone	With heavy metals ions (mean± SD)								
		Ag			Co			Pb		
		5	10	25	5	10	25	5	10	25
C	26.5±0.5	28.5 ±1.5	28 ±2	N.G.	29±1	27.5 ±0.5	28.25±0.25	27 ±3	27.5±0.5	27.5 ±0.5
CN	21.5 ±0.5	20.5 ±0.5	17 ±3	N.G.	20.25±1.75	22 ±0	20.5 ±0.5	22±0	18 ±2	19 ±0
TE	22.5 ±0.5	19.5 ±1.5	23.5 ±0.5	N.G.	23.25 ±1.25	22±1	22.25 ±0.25	20 ±0	23.5 ±0.5	23 ±0
E	R	17 ±1	13.5 ±0.5	N.G.	13 ±3	13.5 ±0.5	12.5 ±0.5	R	12.5 ±2.5	13.75 ±0.75

SD: Standard Deviation

NG: No Growth

R:Resistant

The addition of gentamycin to the medium that supplemented with heavy metals ions showed a significant increasing ($P < 0.01$) in the inhibition zone in the treatment which supplemented with 25 mg / l of silver ions, and no growth was appeared at this treatment , plate (1) .When the addition of tetracycline also a significant increasing ($P < 0.01$) in the inhibition zone and no growth appeared at treatments which containing 25 mg / l of silver ions . On the other hand , the least inhibition zone was found at the treatment which supplemented with 5 mg / l of silver ions , plate (1) and at the treatment with 5 mg / l of lead ions plate (3). When erythromycin used with the supplemented medium with the previous concentrations of heavy metals ions , a significant increasing ($P < 0.01$) was revealed in the inhibition zone at the treatment contain 25 mg / l of silver ions (no growth was appeared) followed by treatment with 5 mg / l of the same metals ions . The inhibition zones were not differ at the other treatment as compared with the control treatment which was free from an additions of heavy metals ions .

In general the inhibition zone in the presence of chloramphenicol was higher significantly ,while leas inhibition zone was appeared with erythromycin when added to the medium supplemented with different heavy metals ions .




Plate (1) : Effect of silver ions with antibiotics on the growth of *P. aeruginosa* on solid medium.




Plate (2): Effect of cobalt ions with antibiotic on the growth of *P. aeruginosa* on solid medium.




Plate (3): Effect of lead ions with antibiotic on the growth of *P. aeruginosa* on solid medium.

The synergistic effect of heavy metals ions and antibiotics on the growth of *S. aureus* in the solid medium

The addition of chloramphenicol antibiotic to the medium which supplemented with 5 or 25 mg/l of silver ions, plate (4) , or cobalt at 25 mg/l , plate (5) or lead at 5 mg/l , plate (6)caused a significant increasing($p<0.01$) in the inhibition zone comparable with the control treatment (free from any addition of heavy metals ions) , table (3).The addition of gentamycin antibiotic to the medium which supplemented with 25 mg/l of silver ions , plate (4) caused a significant increasing($p<0.01$) in the inhibition zone comparable with the control treatment .Also, the addition of vancomycin to the medium which supplemented with different concentration of heavy metals ions showed a significant increasing ($p< 0.01$)in the inhibition zone at the treatment which has 25 mg/l of silver ions , or treatments which have 10 mg/l of cobalt ions ,while the inhibition zone was decreasing largely at the treatment that has 5 mg / l of silver ions, plate (4) .

Table (3):- The inhibition zone diameter (mm) of antibiotic alon and with heavy metals ions on *S. aureus* in the solid medium

Antibiotic	Alone	With heavy metals ions (mean± SD)								
		Ag			Co			Pb		
		5	10	25	5	10	25	5	10	25
C	27.5±0.5	29 ±1	25±1	31±3	23±1	28±4	30±2	33±5	27.5±2.5	26±2
CN	27± 2	27±3	28±4	37±3	23±3	25±1	27.5±0.5	24±0	25±1	22±4
VA	R	R	R	24±4	R	19±1	R	R	R	R
DA	R	29 ± 1	R	40±0	R	30.5±1.5	R	R	R	R

SD: Standard Deviation

R: Resistant




Plate (4) : Effect of silver ions with antibiotic on the growth of *S. aureus* on solid medium .




Plate (5) : Effect of cobalt ions with antibiotic on the growth of *S. aureus* on solid medium .




Plate (6) : Effect of lead ions with antibiotics on the growth of *S. aureus* on solid medium.

The addition of clindamycin antibiotic to the medium which supplemented with different heavy metals ions , a significant increasing ($p < 0.01$) in the inhibition zone was found at the treatment with 5 or 25 mg/ l of silver ions , plate (4) or 10 mg/l of cobalt ions , plate (5) . On the other hand the isolate exhibit a resistant to the antibiotic in the treatment which contains 10 mg /l of silver ions , and 5 or 25 mg/l of cobalt ions , plate (5) ,and all treatments of lead ions , plate (6) .

In general , the inhibition zone of chloramphenicol was the highest , while the vancomycin antibiotic has the the lowest effect as compared with the other antibiotics on the growth of bacteria in the presence of heavy metals ions .

Some *S. aureus* strains resistance to mostly antibiotics (21) . The cause of this resistance beyond to production of some enzymes in bacteria and mis used of antibiotic which lead to multidrugs resistance (22) . Antibiotic resistance of bacteria is increasing worldwide may result public health problems (23) . Multidrugs resistance strains of *P.aeruginosa* are the major cause of mortality in various clinical conditions , such as cystic fibrosis (5) and burn-wound infections (6) , especially in nosocomial situations . This situation has lead to the development of new or modified antimicrobial agents for the therapy of such infection . Heavy metals , particularly silver and mercury , have a variety of applications in controlling microbial

populations (9) . Silver salts alon or in compensations with other drugs appear to have a significant potential as topical antimicrobial agents (13) .

The present study revealed increased suscebtibility of tested isolates toward certin antibiotics when supplemented with heavy metals ions . This results was inline with the results of Higham et al (24) who found that cadmium adapted cells of *Pseudomonas putida* showed greatly increased sensitivity against some antibiotics , including : aminoglycosides , cyclic polypeptides, and doxycyclinen . It is suggested that this is related to changes in outer membrane structure . Also , the results inagreement with results obtained by Ahmed et al (16)

In conclusion heavy metals ions like silver and cobalt alon or in combination with other drugs appear to have a significant potential antimicrobial agents against the major causes of skin infection of human and animals , and burn – wound infections . In addition the heavy metals ions (silver and cobalt) lead to increase the inhibition zones of some antibiotics under study .

تأثير المعادن الثقيلة والمضادات الحياتية على العزلتين الجرثوميتين *Pseudomonas aeruginosa* و *Staphylococcus aureus*

محمد حسن خضر

فرع الإحياء المجهرية – كلية الطب البيطري – جامعة البصرة – البصرة - العراق

الخلاصة

تم تعريض العزلتين الجرثوميتين *Pseudomonas aeruginosa* و *Staphylococcus aureus* الى تراكيز مختلفة (5 ، 10 او 25) ملغم/ لتر من ايونات المعادن الثقيلة (الفضة Ag^{2+} ، الكوبلت Co^{2+} او الرصاص Pb^{2+}) لفترات تعريض مختلفة (0 ، 1 ، 2 ، 24 ، او 48 ساعة).

اوضحت نتائج الدراسة ان التأثير التثبيطي لتلك الايونات على نمو العزلتين كان بالتسلسل التالي

$$Pb < Co < Ag$$

ادت اضافة بعض تلك المعادن الى زيادة الطور التمهيدي lag phase لتلك العزلتين في الوسط الزرعي السائل . و عند اضافة اقراص المضادات الحياتية (*Chloramphenicol* 30 mcg , *Clindamycin* 2 mcg , *Vancomycin* mcg , *Erythromycin* 15 mcg , *Gentamycin* 10 mcg , *Tetracyclin* 30 mcg) الى الاوساط الزرعية الصلبة المدعمة بايونات المعادن بالتراكيز السابقة، اظهرت النتائج حصول تأثير تعاوني بين المضادات الحياتية والمعادن الثقيلة (وبشكل خاص ايونات الفضة بتركيز 25 ملغم / لتر) في زيادة منطقة التثبيط معنويا ($P < 0.01$) لتلك العزلتين .

REFERENCES

1. Groves , D.J. ; Short , J. ; Thewaini , A.J. ; and Young , F.E. (1975) . Epidemiology of antibiotic and heavy metal resistance in bacteria : Resistance patterns in Staphylococci isolated from population in Iraq exposed and not exposed to heavy metals or antibiotics. *Antimicrobiol.Agents and Chemotherapy* , 7(5):622-628.
2. Banerjee , D. and Stablefort , D. (2000) . The treatment of respiratory *Pseudomonas* infection in cystic fibrosis drug and which way ? *Drugs* , 60 : 1053-1064.
3. Conway , S.P. ; Brownlee , K.G.; Denton , M. and Peckham , D.G. (2003) . Antibiotic treatment of multidrug resistant organisms incystic fibrosis . *Am. J. Respir. Med.*, 2 : 321-332.
4. Jones , A.M. ; Govan , J.R. ; Doherty , C.J. ; Dood , M.E. ; Isalska,B.J. ; Stanbridge , T.N. and Webb ,A.K. (2003) .Identification of air borne dissemination of epidemic mutiresistant strains of *Pseudomonas aeruginosa* at a CF centre during a cros infection outbreak . *Thorax* , 58 : 525-527.
5. Canton , R. ; Cobos , N. ; de Gracia ,J. ; Baquero , F. ; Honorato , J.;and Gartner , S. .(2005) . Antimicrobial therapy of pulmonary pathogenic colonization and infection by *Pseudomonas aeruginosa* in cystic fibrosis patients . *Clin. Microbial . Infect.* , 11 : 690- 703.
6. Japoni , A. ; Alborzi , A. ; Kalani , M. ; Nasiri , J. ; Hayati , M. and Farshad ,S. (2006) . Susceptibility patterns and cross-resistance of antibiotics against *Pseudomonas aeruginosa* isolated from burn patients in south of Iran . *Burns* , 32 : 343-347 .
7. Shahid , M. ; Malik , A. and Sheeba (2003) . Multidrug- resistant *Pseudomonas aeruginosa* strains harbouring r-plasmids and a mpc-beta – lactamases isolated from hospitalized burn patients in a tertiary cane hospital of north India . *Fems. Microbiol. Lett.* , 228 :181-186.
8. Shahid , M. and Malik , A.(2005) .Resistance due to aminoglycoside modifying enzymes in *Pseudomonas aeruginosa* isolated from burns patients . *J.Med.Res* . , 122 : 324-329.
9. Kenneth ,A.A. and Jeffrey , T.S.H. (2006) . Manual of dermatologic therapeutics : with essentials of diagnosis . 7th ed. Lippincott Williams and Wilkins , 314pp.

10. Nies , D.H. (1999) . Microbial heavy- metals resistance. Appl. Microbiol. Biotechnol. , 51 : 730-750 . [Medline].
11. Fox,C.L.J.R.; Monafo, W.W.J.R. ; Ayvazian , V.H.; Skinner , A.M. ;Modak,S.; and Stanford , J. (1977) . Topical chemotherapy for burnes using cerium salts and silver sulfadiazine .Surg Gyrecal Obstet , 144 : 668-672.
12. Wassermann,D. ; Schlotterer , M. ; Lebreton , F. ; Levy , J. ; and Guelfi , M. C. (1989) . Use of topically applied silver sulfadiazine plus cerium nitrate in major burns . Burns ,15 : 257-260.
13. De Gracia , C. G. (2001) . An open study comparing topical silver sulfadiazine and topical silver sulfadiazine- cerium nitrate in the treatment of moderate and severe burns. Burns . , 27 : 67-74 .
14. Gerald , E. ; and Mc Donnel (2007) . Antisepsis , disinfection and stterillisation , types action and resistance . Treatment of skin and wound infections . 1st ed. , Blackwell publishing , 156 pp.
15. Chohan , Z.H. ; Munawar , A. ; and Supran , C.T. (2001) .Transition metal ion complexes of Schiff-bases ,synthesis ,characterization and antibacterial properties . Met. Based Drugs , 8 (3) : 137-143 .
16. Ahmed , A. H. ; Omran , A. A.; and El- Sherbiny , G.M. (2006) . Synthesis , characterization and biological evaluation of some methylenedisalicylic acid complexes . J. Appl. Sci . Res . 2 (1) : 44-50.
17. Prasad ,V.S. ; Ballal , M. ; and Shirananda , P.G. (2009) . Action of heavy metals on , *Pseudomonas aerugenosa* strains isolated from infected wounds . J. of Chinese Clinical Medicine , 4(3) :
18. Quinn , P.J. ; Carter , M.E. ; Markey , B.K. ; and Carter , G.R. (1998) . Clinical veterinary microbiology . Mosby , London .
19. Inthorn , D. ; Sidtitoon , N. ; S ilapanuntakul , S. ; and Incharoensakdi , A. (2002) . Sorption of mercury , cadmium , and lead by microalgae . Scince Asia , 28 : 253-261 .
20. Rodriguez, C. E. ; Quesada , A. ; and Rodriguez , E. (2006) . Nickel biosorption by *Acinetobacter baumannii* and *Pseudomonas aerugenosa* isolated from industrial wastewater . Braziliaz J. Microbiol. , 37 : 465- 467 .

21. Gill , S. R. ; Fonts , D.E. ; AR cher , G.L. (2005) . *Staphylococcus aureus* strain and producing methicillin resistant . *Staphylococcus epiderm . Dis .Apr. , 187(7) : 2426-2438 .*
22. Al-Hussainy , D.H. J.M. (2007) . Isolation and diagnosis *Staphylococcus aureus* bacteria from patients infection of urinary tract infection in Al-Diwanyia city . *J . Al-Qadisiya of Vet . Science , 6 (1) : 47-52 .*
23. Maple , P. A. ; Hamilton – Miller , J.L. ; and Barnuffit , W. (1989) . Worldwide antibiotic resistance in methicillin resistant *Staphylococcus aureus* . *Lancet., 2 : 537-540 .*
24. Higham , D.P. ; Sadler , P.J. ; and Scawen , M.D. (1986) . Effect of cadmium on the morphology , membrane integrity and permeability of *Pseudomonas putida* . *J. General Microbiol. , 132 : 1475-1482.*